

Perineal pain after birth: what can help?

Jennifer Hanratty for Evidently Cochrane 9 May 2017 updated 13 January 2021

References

Aasheim V, Nilsen ABV, Lukasse M, Reinart LM. Perineal techniques during the second stage of labour for reducing perineal trauma. Cochrane Database of Systematic Reviews 2011, Issue 12. Art. No.: CD006672. DOI: 10.1002/14651858.CD006672.pub2.

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006672.pub2/abstract>

Plain language summary: http://www.cochrane.org/CD006672/PREG_perineal-techniques-during-the-second-stage-of-labour-for-reducing-perineal-trauma

Abalos E, Sguassero Y, Gyte GML. Paracetamol/acetaminophen (single administration) for perineal pain in the early postpartum period. Cochrane Database of Systematic Reviews 2020, Issue 12. Art. No.: CD008407. DOI: 10.1002/14651858.CD008407.pub3. Accessed 13 January 2021.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD008407.pub3/full>

Plain language summary: http://www.cochrane.org/CD008407/PREG_paracetamol-for-relief-of-perineal-pain-after-birth

Beckmann MM, Stock OM. Antenatal perineal massage for reducing perineal trauma. Cochrane Database of Systematic Reviews 2013, Issue 4. Art. No.: CD005123. DOI: 10.1002/14651858.CD005123.pub3.

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005123.pub3/abstract>

Plain language summary: http://www.cochrane.org/CD005123/PREG_antenatal-perineal-massage-for-reducing-perineal-trauma

East CE, Begg L, Henshall NE, Marchant PR, Wallace K. Local cooling for relieving pain from perineal trauma sustained during childbirth. Cochrane Database of Systematic Reviews 2012, Issue 5. Art. No.: CD006304. DOI: 10.1002/14651858.CD006304.pub3

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006304.pub3/abstract>

Plain language summary: http://www.cochrane.org/CD006304/PREG_local-cooling-for-relieving-pain-from-perineal-trauma-sustained-during-childbirth

Gupta JK, Hofmeyr GJ, Shehmar M. Position in the second stage of labour for women without epidural anaesthesia. Cochrane Database of Systematic Reviews 2012, Issue 5. Art. No.: CD002006. DOI: 10.1002/14651858.CD002006.pub3.

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD002006.pub3/abstract>

Plain language summary: http://www.cochrane.org/CD002006/PREG_position-in-the-second-stage-of-labour-for-women-without-epidural-anaesthesia

Jiang H, Qian X, Carroli G, Garner P. Selective versus routine use of episiotomy for vaginal birth. *Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No.: CD000081. DOI: 10.1002/14651858.CD000081.pub3.

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD000081.pub3/abstract>

Plain language summary: http://www.cochrane.org/CD000081/PREG_selective-versus-routine-use-episiotomy-vaginal-birth

McAndrew, F., Thompson, J., Fellows, L., Large, A., Speed, M., & Renfrew, M. J. (2012). *Infant feeding survey 2010. Leeds: Health and Social Care Information Centre.*

<http://content.digital.nhs.uk/catalogue/PUB08694/Infant-Feeding-Survey-2010-Consolidated-Report.pdf>

Molakatalla S, Shepherd E, Grivell RM. Aspirin (single dose) for perineal pain in the early postpartum period. *Cochrane Database of Systematic Reviews* 2017, Issue 2. Art. No.: CD012129. DOI: 10.1002/14651858.CD012129.pub2.

<http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD012129.pub2/abstract>

Plain language summary: http://www.cochrane.org/CD012129/PREG_aspirin-single-dose-relief-perineal-pain-after-childbirth

Wuytack F, Smith V, Cleary BJ. Oral non-steroidal anti-inflammatory drugs (single dose) for perineal pain in the early postpartum period. *Cochrane Database of Systematic Reviews* 2021, Issue 1. Art. No.: CD011352. DOI: 10.1002/14651858.CD011352.pub3. Accessed 13 January 2021.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD011352.pub3/full>

Plain language summary: http://www.cochrane.org/CD011352/PREG_anti-inflammatory-drugs-relief-perineal-pain-after-childbirth

UNICEF. *Caring for your baby at night A guide for parents* NCT.org: UNICEF UK Baby Friendly Initiative.