

Traumatic brain injury: is decompressive surgery worth the risks?

Jason Yuen for Evidently Cochrane 3 April 2020

References

- [1] Friedland D, Hutchinson P. Classification of traumatic brain injury. *Advances in Clinical Neuroscience and Rehabilitation* 2013. Available from: <https://www.acnr.co.uk/2013/07/classification-of-traumatic-brain-injury/>
- [2] Marsh H. *Do No Harm: Stories of Life, Death and Brain Surgery*. London: Weidenfeld & Nicolson, 2014. <https://www.amazon.co.uk/Do-No-Harm-Stories-Surgery/dp/0297869876>
- [3] Sahuquillo J, Dennis JA. Decompressive craniectomy for the treatment of high intracranial pressure in closed traumatic brain injury. *Cochrane Database of Systematic Reviews* 2019, Issue 12. Art. No.: CD003983. DOI: 10.1002/14651858.CD003983.pub3. <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD003983.pub3/full>
- [4] Higgins JPT, Altman DG, Sterne JAC (editors). Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. https://handbook-5-1.cochrane.org/chapter_8/box_8_11_a_a_note_on_blinding_in_clinical_trials.htm